

## Phylogenetic Analysis of RhoGAP Domain-Containing Proteins

Marcelo M. Brandão<sup>1</sup>, Karina L. Silva-Brandão<sup>2</sup>, Fernando F. Costa<sup>1</sup>, and Sara T.O. Saad<sup>1\*</sup>

<sup>1</sup>*Hematology and Hemotherapy Center, State University of Campinas, 13083-970 Campinas, SP, Brazil;*

<sup>2</sup>*Department of Zoology, Institute of Biology, State University of Campinas, 13083-970 Campinas, SP, Brazil.*

**Proteins containing an Rho GTPase-activating protein (RhoGAP) domain work as molecular switches involved in the regulation of diverse cellular functions. The ability of these GTPases to regulate a wide number of cellular processes comes from their interactions with multiple effectors and inhibitors, including the RhoGAP family, which stimulates their intrinsic GTPase activity. Here, a phylogenetic approach was applied to study the evolutionary relationship among 59 RhoGAP domain-containing proteins. The sequences were aligned by their RhoGAP domains and the phylogenetic hypotheses were generated using Maximum Parsimony and Bayesian analyses. The character tracing of two traits, GTPase activity and presence of other domains, indicated a significant phylogenetic signal for both of them.**

**Key words:** Bayesian analysis, character tracing, parsimony, phylogenomics, protein domain, RhoGAP

### Introduction

The Rho GTPase-activating proteins (RhoGAPs) are defined by the presence of a 150-amino-acid homolog region that is designated as the RhoGAP domain. This domain is necessary and sufficient for GAP activity and shares at least 20% sequence identity among its family members (1, 2). Proteins containing an RhoGAP domain act as molecular switches involved in the regulation of diverse cellular functions, including actin cytoskeleton rearrangements, regulation of gene transcriptions, cell cycle regulation, control of apoptosis, and membrane trafficking (2–5). Rho GTPases cycle between active and inactive GTP-bound states. The control of these states is regulated by three main classes of proteins: guanine nucleotide exchange factors, guanine nucleotide dissociation inhibitors, and GAPs. To date, at least 21 Rho GTPases have been defined, among which only three (RhoA, Cdc42, and Rac1) are well characterized. Therefore, most studies have been focusing on these three proteins.

The ability of Rho GTPases to regulate a wide number of cellular processes comes from their interactions with multiple effectors or inhibitors. One class of these inhibitors is the RhoGAP family, which stimulates GTPase activity by enhancing the intrinsic rate

of GTP hydrolysis.

In the early analyses of the human genome sequence, 77 different genes containing the RhoGAP domain were found. Further studies have demonstrated that many of these genes are simple gene sequence variations or single nucleotide polymorphisms (6, 7). The structural data available for RhoGAP domain-containing proteins showing their complexity with Rho GTPases (Cdc42 and RhoA) demonstrated the 3D workflow for RhoGAP-mediated GTP-hydrolysis, and highlighted the importance of a well-conserved arginine residue present in the active site that acts as a conformation stabilizer needed for hydrolysis (8–10).

Recently, a novel member of the RhoGAP family, ARHGAP21 (Rho GTPase-activating protein 21, alias ARHGAP10), was cloned and characterized in our laboratory (11). In addition to the RhoGAP domain, ARHGAP21 presents a PH domain and a P-loop-containing PDZ domain. This gene is widely expressed at high levels in muscle and brain, and is up-regulated during myeloid and erythroid maturation, suggesting a potential role for this RhoGAP in regulating cell differentiation (11).

The aim of this study is to infer the evolution of the RhoGAP superfamily using a phylogenetic approach, to determine the roles of other domains and their main GTPase activities in this evolutionary his-

**\*Corresponding author.**

**E-mail:** sara@unicamp.br

tory, and to provide a tool that could render some insights regarding subfamily protein functions using RhoGAP-containing proteins as a model.

## Results

The full dataset contains 267 amino acids, of which 161 are parsimony-informative. Parsimony searches of the equally weighed dataset resulted in 12 equally parsimonious trees with 3,213 steps (CI=0.449; RI=0.525). The strict consensus tree was mostly not resolved at internal nodes (data not shown), but almost all terminal branches showed strong bootstrap support (Figure 1).

A Bayesian tree recovered mostly the same terminal relationships by Maximum Parsimony (MP) analyses with strong values of Posterior Probability (PP). However, internal branches have from low to moderate PP values (Figure 1).

The two characters investigated (domains and GTPase activity) showed a significant phylogenetic signal ( $P=0.003$ ), which suggests that the distribution of these traits among the proteins can be explained by their phylogenetic relationships (Figure 2). The optimization of the other domains over our phylogenetic hypothesis suggests that the ancestral state of these proteins involves solely the presence of the RhoGAP domain and the activity toward Rac1.

The character tracing of these traits suggests an overall pattern on which proteins sharing equal domains also share equal GTPase activity. The clade joining KIAA0672 + 3BP1 + RICH1 + Nadrin was recovered both by MP and Bayesian analyses with strong support. All proteins in this clade share the presence of the BAR (Bin-Amphiphysin-Rvs) domain and GTPase activity toward Rac1, solely or in addition to other GTPases. The same rule can be applied to the clade joining STARTdom + GT650 + DLC1 + AHRGAP7. All of them share the presence of the START (steroidogenic acute regulatory protein-related lipid transfer) domain and all, but STARTdom, have GTPase activity over RhoA. GTPase activity is unknown for STARTdom. The clade joining GMIP + HA1 + PARG is composed by proteins containing the C1 domain in addition to their GTPase activity toward RhoA.

The clade joining AHRGAP11A + AHRGAP20 + AHRGAP1 + AHRGAP8 has in common the absence of other domains that are not RhoGAP. The GTPase activity of this group is known only for AHRGAP1, which is active toward Cdc42. Considering other ter-

minal clades, we can presume that other AHRGAPs within this group can show the same GTPase activity toward Cdc42.

On the other hand, the clade joining P115 + srGAP3 + srGAP1 + srGAP2 shares the presence of the FCH (FER/CIP4-homology) domain, however, P115 has GTPase activity over Rac1, while the other three proteins have activity toward Cdc42.


## Discussion

Phylogenetic reconstruction and bioinformatics analyses that integrate evolutionary considerations are becoming increasingly important tools for applied fields. Numerous gene sequences were generated in the genomics age with little or no accompanying experimental determination of functional information or evolutionary relationships. Previous works from Peck *et al* (12) and Moon and Zheng (2) also present a phylogenetic approach on the RhoGAP family; however, the authors did not indicate the methodology applied neither did they present any support analyses for their cladograms.


In this work, bioinformatics and phylogenomics tools were used to present a phylogenetic relationship of 59 members of the RhoGAP superfamily. All amino acid alignments and subsequent phylogenetic tree constructions were based on the RhoGAP domain sequence. We demonstrated that these RhoGAP domain-containing proteins, with the conservative arginine residue, form a monophyletic group, that is, all of them share a common protein ancestor in their evolutionary history.

The tracing for GAP activity toward the most studied RhoGTPases (RhoA, Rac1, and Cdc42) (Figure 2) indicates that this trait presents a strong phylogenetic signal ( $P=0.003$ ), contrasting with previous findings of Peck *et al* (12).

The analysis of the resulting phylogenetic tree has suggested that the ancestral state for GTPase activity is the affinity to Rac1. It is still difficult to determine the gap activity by only analyzing the protein sequence; the GAP assay (13, 14) is the most reliable way to determine activity. The phylogenetic approach may give a clue, once it is capable of clustering together different proteins that share common substrates as can be seen on the clades of KIAA0672 + 3BP1 + RICH1 + Nadrin and srGAP3 + srGAP1 + srGAP2. Speculations regarding protein specific functions (only using the GTPase activity character)



**Fig. 1** The Bayesian tree based on amino acid sequences of the RhoGAP domain from RhoGAP domain-containing proteins. Values above the branches indicate the Posterior Probability (PP) and the bold numbers indicate the bootstrap values from 500 replications (where it exceeds 50%).



**Fig. 2** Domain presence (left side) and GTPase activity (right side) in the RhoGAP domain-containing proteins traced along the proposed phylogeny.

may be avoided for now, because the affinity for the same GTPase does not imply in the same function, since each GTPase may present contrasting functions in different pathways (15).

Furthermore, structural and molecular biology studies are needed to elucidate the exact amino acid composition involved in determining specificity and how the differences in this composition can affect the 3D protein structure and its interaction with Rho GTPases.

In addition to the RhoGAP domain, the members of this superfamily usually contain other functional motifs. Therefore, RhoGAPs might catalyze or participate in enzymatic reactions other than the enhancement of the intrinsic GTP hydrolysis of Rho GTPases, and sometimes apparently aiding the Rho protein to signaling (2).

Somehow the presence of additional domains was linked to the RhoGAP domain structure because, even focusing the alignment exclusively on the RhoGAP domain sequence, the phylogeny joined in clades of different proteins sharing the same additional domains with strong bootstrap and PP, that is, the ARHGAPs were divided into two groups. One is composed of a clade including the ARHGAPs 9, 12, 21, and 23, presenting an RhoGAP domain and a PH domain accompanied or not by additional domains (Figure 2). The other is composed of the terminals ARHGAPs 1, 8, 11A, and 20, presenting only the RhoGAP domain, except the ARHGAP20 that present an RA domain.

Interactions among genomics, evolution, and bioinformatics go further than sequence alignment and relationship elucidation among species. Evolutionary analysis may help researchers design new strategies to understand protein or gene interactions and their functionalities and might provide an insight for new experiments. In conclusion, a phylogenetic study of the RhoGAP domain-containing proteins has demonstrated that there is a strong evolutionary relationship among the RhoGAP superfamily members, especially when they share common motifs or GAP activity.

## Materials and Methods

### Materials

All protein sequences used here were obtained from the GenBank database (<http://ncbi.nlm.nih.gov/Genbank/>) at the National Center for Biotechnol-

ogy Information (NCBI), as well as from the Swiss-Prot/TrEMBL database (<http://expasy.org/sprot/>) at the Swiss Institute for Bioinformatics and at the European Bioinformatics Institute (Table 1).

The sequences were aligned by their RhoGAP domains with additional 100 N-terminal residues, primarily using ClustalW version 1.83 (16) under default settings, followed by adjustment by eye using the BioEdit version 6.0.7 (Ibis Therapeutics, Carlsbad, USA). All alignment files, the protein sequences in the FASTA format, and other related colored materials are available for download at <http://www.hemocentro.unicamp.br/submission/>.

### Phylogenetic analyses

The Bayesian analysis was carried out by using Mr-Bayes, version 3.1.2 (17, 18) with the mixed model of amino acid substitution provided in the package. Six simultaneous chains were conducted for  $1.0 \times 10^6$  generations, sampling trees every 100 cycles. The first 1,000 trees were discarded as "burn in". For all analyses, chr5orf was used as an outgroup to root the tree, based on the absence of the conservative argentine residue.

The MP analyses were performed with PAUP\* 4.0b10 (19) on the entire dataset using a heuristic search with 500 random taxon addition replicates, TBR, branch-swapping, gaps scored as missing data, and all characters equally weighted. A strict consensus tree was computed whenever multiple equally parsimonious trees were obtained. The robustness of each branch was determined using the nonparametric bootstrap test (20) with 500 replicates and 10 random taxon additions.

### Character optimization

MacClade 4.08 (21) was used to perform character optimization analyses. We investigated the evolution of two characters that were superimposed onto the Bayesian tree proposed for the RhoGAP-containing proteins: the presence of different domains in addition to RhoGAP, and the GTPase enhancing activity toward the most studied Rho GTPases (Rac1, RhoA, and Cdc42). For domain identification, a search of the PFAM database version 19.0 (22) was performed using the HMMPFAM tool from the HMMER suite version 2.3.2 (23) with the E-value cutoff for the per-sequence set to  $1.0E-10$ . This character had 20 unordered character states plus a 21<sup>st</sup> character state

Table 1 Selected Rho GTPase-Activating Proteins

No.	Protein	GenBank	SwissProt	No.	Protein	GenBank	SwissProt
1	3BP-1		Q9Y3L3	31	HA-1 (KIAA0233)	BAA13212.1	
2	ABR	NP_068781.2		32	H-Graf	CAA71414.2	
3	$\alpha$ -Chimaerin	CAA35769.1		33	INPP5B	AAA79207.1	
4	ARAP1	NP_056057.1		34	KIAA0672	BAA31647	
5	ARAP2	BAA25506.1		35	KIAA1204	BAA86518.1	
6	ARAP3	CAC83946.1		36	KIAA1314	BAA92552.1	
7	ARHGAP1	NP_004299		37	KIAA1688	BAB21779.1	
8	ARHGAP6	NP_038286.1		38	MgcRacGAP	NP_037409.2	
9	ARHGAP7		Q63744	39	Myosin_IXA	NP_008832.1	
10	ARHGAP8	CAB90248.1		40	Myosin_IXB	NP_004136.2	
11	ARHGAP9	BAB56159.1		41	Nadrin	NP_060524.4	
12	ARHGAP11A	NP_055598.1		42	N-Chimaerin homolog	AAB81198.1	
13	ARHGAP12	NP_060757.4		43	OCRL-1	NP_001578.2	
14	ARHGAP18 MacGAP	NP_277050		44	Oligophrenin-1	NP_002538.1	
15	ARHGAP19	NP_116289.4		45	p190-A	AAF80386.1	
16	ARHGAP20	AAS45466.1		46	p190-B	NP_001164.2	
17	ARHGAP21	AF480466.1		47	P85-alpha		P27986
18	ARHGAP23	BAA96025.1		48	P85-beta	NP_005018.1	
19	ARHGAP24	NP_112595.1		49	PARG1	NP_004806.2	
20	ARHGAP25b	NP_055697.1		50	PSGAP	AAK18175.1	
21	$\beta$ -Chimaerin	AAA16836.1		51	RALBP1	NP_006779.1	
22	BCR	NP_004318.2		52	RHG4 (p115)	CAA55394.1	
23	CAC17688.2	CAC17688.2		53	RICH-1	CAC37948.1	
24	CHR5ORF	NP_057687.1		54	RLIP76	AAB00103.1	
25	DLC-1	NP_006085.2		55	srGAP1	BAA92542.1	
26	GAPDro	AAF44627.1		56	srGAP2	BAA32301.1	
27	GMIP	NP_057657.1		57	srGAP3	CAC22407.1	
28	GRAF	NP_055886.1		58	START domain	NP_055540.2	
29	GRAF-2	BAB61771			containing 8 (KIAA0189)	CAC22407.1	
30	GT650 (DLC2)	NP_443083.1		59	HA-1 (KIAA0233)	BAA13212.1	

corresponding to the absence of additional domains other than RhoGAP. The GTPase character had eight character states representing the affinity toward one or two Rho GTPases; these data was mined by searching PubMed (<http://www.ncbi.nlm.nih.gov/entrez>) at NCBI.

To test whether there was a phylogenetic signal in the characters traced, we used the methodology proposed by Wahlberg (24) that was modified from the PTP test described by Faith and Cranston (25). The test compared the number of steps of the tree constructed with the actual data, with the number of steps in the trees obtained for each random reshuffling of the separated character states. We performed 300 random reshufflings of character states among the fixed terminal proteins by using Mesquite version 1.06

(<http://www.mesquiteproject.org>). The probability ( $P$ ) that the observed pattern does not differ from random is given as  $(n + 1)/300$ , where  $n$  is the number of replications no bigger than that of the actual steps. A significant phylogenetic signal was observed when  $P$  is less than 0.05 (25).

## Acknowledgements

This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP; Grant 03/06621-5 to MMB) and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES; fellowship to KLSB). We thank Ms. Raquel Foglio for English revision.

### Authors' contributions

MMB participated in the design of the study, sequence alignment, bioinformatics analyses, and drafted the manuscript. KLSB participated in all automatic alignment and eye refinements, interpretations of the bioinformatics results, and drafted the manuscript. FFC and STOS conceived the study, participated in its design and coordination, and helped to draft the manuscript. All authors read and approved the final manuscript.

### Competing interests

The authors have declared that no competing interests exist.

### References

- Aitsebaomo, J., *et al.* 2004. p68RacGAP is a novel GTPase-activating protein that interacts with vascular endothelial zinc finger-1 and modulates endothelial cell capillary formation. *J. Biol. Chem.* 279: 17963-17972.
- Moon, S.Y. and Zheng, Y. 2003. Rho GTPase-activating proteins in cell regulation. *Trends Cell Biol.* 13: 13-22.
- Hall, A. 1998. Rho GTPases and the actin cytoskeleton. *Science* 279: 509-514.
- Lamarche, N. and Hall, A. 1994. GAPs for rho-related GTPases. *Trends Genet.* 10: 436-440.
- Van Aelst, L. and D'Souza-Schorey, C. 1997. Rho GTPases and signaling networks. *Genes Dev.* 11: 2295-2322.
- Sachidanandam, R., *et al.* 2001. A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms. *Nature* 409: 928-933.
- Lander, E.S., *et al.* 2001. Initial sequencing and analysis of the human genome. *Nature* 409: 860-921.
- Nassar, N., *et al.* 1998. Structures of Cdc42 bound to the active and catalytically compromised forms of Cdc42GAP. *Nat. Struct. Biol.* 5: 1047-1052.
- Rittinger, K., *et al.* 1997. Crystal structure of a small G protein in complex with the GTPase-activating protein rhoGAP. *Nature* 388: 693-697.
- Rittinger, K., *et al.* 1997. Structure at 1.65 Å of RhoA and its GTPase-activating protein in complex with a transition-state analogue. *Nature* 389: 758-762.
- Basseres, D.S., *et al.* 2002. ARHGAP10, a novel human gene coding for a potentially cytoskeletal Rho-GTPase activating protein. *Biochem. Biophys. Res. Commun.* 294: 579-585.
- Peck, J., *et al.* 2002. Human RhoGAP domain-containing proteins: structure, function and evolutionary relationships. *FEBS Lett.* 528: 27-34.
- Manser, E., *et al.* 1995. Identification of GTPase-activating proteins by nitrocellulose overlay assay. *Methods Enzymol.* 256: 130-139.
- Manser, E., *et al.* 1992. Diversity and versatility of GTPase activating proteins for the p21rho subfamily of ras G proteins detected by a novel overlay assay. *J. Biol. Chem.* 267: 16025-16028.
- Bishop, A.L. and Hall, A. 2000. Rho GTPases and their effector proteins. *Biochem. J.* 348: 241-255.
- Thompson, J.D., *et al.* 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22: 4673-4680.
- Ronquist, F. and Huelsenbeck, J.P. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572-1574.
- Huelsenbeck, J.P. and Ronquist, F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754-755.
- Swofford, D.L. 2002. PAUP\*: phylogenetic analysis using parsimony (and other methods), version 4.0b10. CD-ROM. Sinauer Associates, Sunderland, USA.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783-791.
- Maddison, W.P. and Maddison, D.R. 1999. MacClade: analysis of phylogeny and character evolution, version 3.08. CD-ROM. Sinauer Associates, Sunderland, USA.
- Sonnhammer, E.L., *et al.* 1997. Pfam: a comprehensive database of protein domain families based on seed alignments. *Proteins* 28: 405-420.
- Eddy, S.R. 1998. Profile hidden Markov models. *Bioinformatics* 14: 755-763.
- Wahlberg, N. 2001. The phylogenetics and biochemistry of host-plant specialization in Melitaeine butterflies (Lepidoptera: Nymphalidae). *Evolution* 55: 522-537.
- Faith, D.P. and Cranston, P.S. 1991. Could a cladogram this short have arisen by chance alone? On permutation tests for cladistic structure. *Cladistics* 7: 1-28.