

## **Genomics Proteomics Bioinformatics**



www.elsevier.com/locate/gpb www.sciencedirect.com

## **CORRIGENDUM**

## Corrigendum to "Antibiotic Treatment Drives the Diversification of the Human Gut Resistome" [Genomics Proteomics Bioinformatics 17 (1) (2019) 39-51]



Jun Li $^{1,2,\#,a}$ , Elizabeth A. Rettedal $^{3,\#,b}$ , Eric van der Helm $^{3,c}$ , Mostafa Ellabaan $^{3,d}$  Gianni Panagiotou $^{4,5,6,*,e}$ , Morten O.A. Sommer $^{3,*,f}$ 

Available online 9 September 2020

The editors regret there was an error in "Shotgun metagenome library construction and sequencing" section. "The raw sequences can be found in BGID (CRA000815)" should be corrected to "The raw metagenome sequencing data have been deposited in the Genome Sequence Archive at Beijing Institute of Genomics, Chinese Academy of Sciences (GSA: CRA000815), and are publicly accessible at https://bigd.big. ac.cn/gsa/". The correct section is shown below. The editors would like to apologize for any inconvenience

DOI of original article: 10.1016/j.gpb.2018.12.003.

Peer review under responsibility of Beijing Institute of Genomics, Chinese Academy of Sciences and Genetics Society of China.

<sup>&</sup>lt;sup>1</sup> Department of Infectious Diseases and Public Health, Colleague of Veterinary Medicine and Life Sciences, City University of Hong Kong, Hong Kong Special Administrative Region, China

<sup>&</sup>lt;sup>2</sup> School of Data Science, City University of Hong Kong, Hong Kong Special Administrative Region, China

<sup>&</sup>lt;sup>3</sup> Novo Nordisk Foundation Center for Biosustainability, DK-2900 Hørsholm, Denmark

<sup>&</sup>lt;sup>4</sup> Systems Biology and Bioinformatics Unit, Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute, 07745 Jena, Germany

<sup>&</sup>lt;sup>5</sup> Systems Biology and Bioinformatics Group, School of Biological Sciences, Faculty of Sciences, The University of Hong Kong, Hong Kong Special Administrative Region, China

<sup>&</sup>lt;sup>6</sup> Department of Microbiology, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong Special Administrative Region, China

Corresponding authors.

E-mail: gianni.panagiotou@leibniz-hki.de (Panagiotou G), msom@bio.dtu.dk (Sommer MOA).

<sup>#</sup> Equal contribution.

<sup>&</sup>lt;sup>a</sup> ORCID: 0000-0001-7218-429X.

b ORCID: 0000-0001-5586-0508.

c ORCID: 0000-0002-1669-2529.

<sup>&</sup>lt;sup>d</sup> ORCID: 0000-0002-9736-0461.

e ORCID: 0000-0001-9393-124X.

f ORCID: 0000-0003-4005-5674.

## Shotgun metagenome library construction and sequencing

Culture-independent fecal extracts from treated subject samples Day 0, Day 2, Day 5, Day 12, Day 19, Day 26, Day 33, and Day 97 and control subject samples Day 0, Day 2, Day 19, and Day 97 were used to build shotgun metagenomic

libraries using the Nextera XT kit with the standard protocol. The HiSeq 1500 was used for 100 bp PE sequencing in the CGS of The University of Hong Kong and the average throughput for each sample was 10.5 Gbp. The raw metagenome sequencing data have been deposited in the Genome Sequence Archive at Beijing Institute of Genomics, Chinese Academy of Sciences (GSA: CRA000815), and are publicly accessible at https://bigd.big.ac.cn/gsa/.