

Question of the Animal-Plant-Fungal Divergence

Sachiko Matsutani

Division of Microbiology, National Institute of Health Sciences, Japan.

There has been a controversial issue in phylogenetic relationships among animals, fungi, and plants. Before Whittaker and Margulis (1) classified the fungi as a separate kingdom in their five-kingdom classification, fungi traditionally had been considered more closely related to plants than to animals. With the determination of the primary structures of small subunit rRNA and the proteins like actin and α -tubulin in various organisms, these sequences have been used to make molecular trees. Most of these studies show that fungi and animals are sister groups (2–4).

More recent investigations of the proteins involved in RNA metabolism, the mRNA capping apparatus, and several key components that regulate the cell cycle, suggest a close relationship between animals and plants, with fungi as more distant (5–7). Moreover, there are intriguing examples of gene families and domain structures shared exclusively by animals and plants (8–10), as referred by Stiller in this issue (11). Phylogenetic analyses of several additional proteins also show that there is a tree building signal that clusters animal and plant sequences (11). Based on the recent data, now Stiller (11) reviews new lines of evidence that show a close relationship between animals and plants, and points out that these lines of evidence have to be incorporated into models of eukaryotic evolution in the future. Certainly, recent bioinformatic, proteomic, and genomic data have great potential for helping to resolve ancient evolutionary relationships. These data will provide further understanding of the relationships among animals, fungi, and plants, like accumulations of the sequence data of rRNA and ubiquitous proteins had ever resulted in hypotheses, but now on the broader scales.

*** Corresponding author.**

E-mail: sachiko@nihs.go.jp

References

1. Whittaker, R.H. and Margulis, L. 1978. Protist classification and the kingdoms of organisms. *Biosystems* 10: 3-18.
2. Baldauf, S.L. and Palmer, J.D. 1993. Animals and fungi are each other's closest relatives: congruent evidence from multiple proteins. *Proc. Natl. Acad. Sci. USA* 90: 11558-11562.
3. Wainright, P.O., *et al.* 1993. Monophyletic origins of the metazoa: an evolutionary link with fungi. *Science* 260: 340-342.
4. Borchiellini, C., *et al.* 1998. Phylogenetic analysis of the Hsp70 sequences reveals the monophyly of Metazoa and specific phylogenetic relationships between animals and fungi. *Mol. Biol. Evol.* 15: 647-655.
5. Anantharaman, V., *et al.* 2002. Comparative genomics and evolution of proteins involved in RNA metabolism. *Nucleic Acids Res.* 30: 1427-1464.
6. Shuman, S. 2002. What messenger RNA capping tells us about eukaryotic evolution. *Nature Reviews Mol. Cell Biol.* 3: 619-625.
7. Oakenfull, E.A., *et al.* 2002. Plant D-type cyclins and the control of G1 progression. *Philos. Trans. R. Soc. Lond., B* 357: 749-760.
8. Pandey, R., *et al.* 2002. Analysis of histone acetyltransferase and histone deacetylase families of *Arabidopsis thaliana* suggests functional diversification of chromatin modification among multicellular eukaryotes. *Nucleic Acids Res.* 30: 5036-5055.
9. Verelst, W. and Asard, H. 2003. A phylogenetic study of cytochrome b561 proteins. *Genome Biol.* 4: R38.
10. Matsutani, S. 2004. Similarities in transcription factor IIC subunits that bind to the posterior regions of internal promoters for RNA polymerase III. *BMC Evol. Biol.* 4: 26.
11. Stiller, J.W. 2004. Emerging genomic and proteomic evidence on relationships among the animal, plant and fungal kingdoms. *Geno. Prot. Bioinfo.* 2: 70-76.